



Product Name: M+PAC MYCOPLASMA HYOPNEUMONIAE INACTIVATED VACCINE FOR PIGS
APVMA Approval No: 62077/138481

Label Name:	M+PAC MYCOPLASMA HYOPNEUMONIAE INACTIVATED VACCINE FOR PIGS
Signal Headings:	FOR ANIMAL TREATMENT ONLY
Constituent Statements:	<p>A liquid adjuvanted bacterin consisting of ≥ 1.1 RP units inactivated Mycoplasma hyopneumoniae strain J.</p> <p>Also contains: 0.134 mL mineral oil (adjuvant), 0.096 mL aluminium hydroxide (adjuvant), 0.01 mg/mL thiomersal (preservative)</p>
Claims:	For the vaccination of healthy pigs as an aid in the prevention of pneumonia caused by Mycoplasma hyopneumoniae infection.
Net Contents:	<p>100 mL (50 x 2 mL doses or 100 x 1 mL doses) 250 mL (125 x 2 mL doses or 250 x 1 mL doses) 500 mL (250 x 2 mL doses or 500 x 1 mL doses)</p>
Directions for Use:	
Restrictions:	
Contraindications:	This vaccine is not to be used on breeder stock.
Precautions:	Use with caution as compatibility studies with concurrent use of other veterinary products were not performed.
Side Effects:	Transient local reaction may occur at the injection site.
Dosage and Administration:	<p>Use the contents immediately after broaching the vial/container. Discard the unused portion.</p> <p>Shake well before use.</p>

	<p>Using aseptic technique vaccinate piglets.</p> <p>7-10 days of age or older: 1 mL dose subcutaneously or intramuscularly followed by a second 1 mL dose 14 days later.</p> <p>3 weeks of age or older (for herds using a single dose program): 2 mL intramuscularly. Intramuscular injection with a single 2 mL dose at 3 weeks of age has been demonstrated to provide at least 24 weeks duration of immunity.</p>
--	--

General Directions:	<p>CAUTION: AVOID CARCASS DAMAGE</p> <ol style="list-style-type: none"> 1. Sterilise all injection apparatus by boiling before use. Avoid use of strong disinfectants on apparatus. 2. Maintain maximum cleanliness at all times. 3. Keep needles sharp and clean. Use needles of appropriate gauge and length. 4. Check that the equipment is functioning properly and that it is delivering the correct dose. 5. As far as possible avoid injection of animals in wet weather or under dusty conditions. 6. Inject high on the side of the neck behind the ear.
---------------------	--

Withholding Periods:	Zero (0) days.
----------------------	----------------

Trade Advice:	EXPORT SLAUGHTER INTERVAL (ESI): Zero (0) days.
---------------	---

Safety Directions:	
--------------------	--

First Aid Instructions:	If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If swallowed, do NOT induce vomiting.
-------------------------	--

First Aid Warnings:	
---------------------	--

Additional User Safety:	<p>Take care to avoid self-injection.</p> <p>This product contains mineral oil and is an irritant. In the event of self-administration, it can cause significant pain and swelling at the injection site, perhaps also involving the draining lymph nodes. Medical or surgical intervention may be required. Contact a doctor immediately, even if only a very small amount is injected, and take this package leaflet/ carton with you. Allow the wound to bleed freely and DO NOT squeeze or interfere with the injection site.</p> <p>Ancillary advice to the medical practitioner This product contains mineral oil. Even small amounts of self-administered mineral oil can cause intense swelling and a persistent granulomatous inflammatory reaction. If injected into a finger joint or tendon sheath, the product may track along the tendon, perhaps also involving the draining lymph nodes. The swelling and inflammation may compromise blood supply and result in necrosis that, in rare cases, may lead to the loss of a digit.</p> <p>Following appropriate immediate local cleansing, corticosteroids may be considered to decrease the severity of any local reaction. Ascertain the patient's tetanus immunisation status, and provide booster or primary series, as appropriate.</p> <p>In some cases of self-injection, PROMPT surgical attention may be required. The wound should be incised and irrigated to remove the vaccine, especially where there is involvement of finger pulp or tendon. Complete curettage or total excision of the lesion may</p>
-------------------------	--

be required for chronic granulomatous reactions. Meticulous technique is required to stop inadvertent spread of the product.

Environmental
Statements:

Disposal:

Dispose of containers and outer packaging by wrapping with paper and putting in garbage. Discarded needles should immediately be placed in a designated and appropriately labelled 'sharps' container.

Storage:

Store between 2 °C and 8 °C (refrigerate, do not freeze). Protect from light. Keep out of reach of children.